



# THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### On $\mathbb{Q}$ -de Rham cohomology via $\mathbb{Q}$ -rings

#### Citation for published version:

Pridham, JP 2016 'On  $\mathbb{Q}$ -de Rham cohomology via  $\mathbb{Q}$ -rings' ArXiv.

#### Link:

[Link to publication record in Edinburgh Research Explorer](#)

#### General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

#### Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



ON  $q$ -DE RHAM COHOMOLOGY VIA  $\Lambda$ -RINGS

J.P. PRIDHAM

ABSTRACT. We show that Aomoto's  $q$ -deformation of de Rham cohomology arises as a natural cohomology theory for  $\Lambda$ -rings. Moreover, Scholze's  $(q-1)$ -adic completion of  $q$ -de Rham cohomology depends only on the Adams operations at each residue characteristic. This gives a fully functorial cohomology theory, including a lift of the Cartier isomorphism, for smooth formal schemes in mixed characteristic equipped with a suitable lift of Frobenius. If we attach  $p$ -power roots of  $q$ , the resulting theory is independent even of these lifts of Frobenius, refining a comparison by Bhatt, Morrow and Scholze.

## INTRODUCTION

The  $q$ -de Rham cohomology of a polynomial ring is a  $\mathbb{Z}[q]$ -linear complex given by replacing the usual derivative with the Jackson  $q$ -derivative  $\nabla_q(x^n) = [n]_q x^{n-1} dx$ , where  $[n]_q$  is Gauss'  $q$ -analogue  $\frac{q^n-1}{q-1}$  of the integer  $n$ . In [Sch2], Scholze discussed the  $(q-1)$ -adic completion of this theory for smooth rings, explaining relations to  $p$ -adic Hodge theory and singular cohomology, and conjecturing that it is independent of co-ordinates.

We show that  $q$ -de Rham cohomology naturally arises as a functorial invariant of  $\Lambda$ -rings (Theorem 1.11), and that its  $(q-1)$ -adic completion depends only on a  $\Lambda_P$ -ring structure (Theorem 2.7), for  $P$  the set of residue characteristics; a  $\Lambda_P$  ring has a lift of Frobenius for each  $p \in P$ . This recovers the known equivalence between de Rham cohomology and complete  $q$ -de Rham cohomology over the rationals, while giving no really new functoriality statements for smooth schemes over  $\mathbb{Z}$ . However, in mixed characteristic, it means that complete  $q$ -de Rham cohomology depends only on a lift  $\Psi^p$  of absolute Frobenius locally generated by co-ordinates with  $\Psi^p(x_i) = x_i^p$ . Given such data, we construct (Proposition 2.8) a quasi-isomorphism between Hodge cohomology and  $q$ -de Rham cohomology modulo  $[p]_q$ , extending the local lift of the Cartier isomorphism in [Sch2, Proposition 3.4].

Taking the Frobenius stabilisation of the complete  $q$ -de Rham complex of  $A$  yields a complex resembling the de Rham–Witt complex. We show (Theorem 3.10) that up to  $(q^{1/p^\infty} - 1)$ -torsion, the  $p$ -adic completion of this complex depends only on the  $p$ -adic completion of  $A[\zeta_{p^\infty}]$  (where  $\zeta_n$  denotes a primitive  $n$ th root of unity), with no requirement for a lift of Frobenius or a choice of co-ordinates. The main idea is to show that the stabilised  $q$ -de Rham complex is in a sense given by applying Fontaine's period ring construction  $A_{\text{inf}}$  to the best possible perfectoid approximation to  $A[\zeta_{p^\infty}]$ . As a consequence, this shows (Corollary 3.11) that after attaching all  $p$ -power roots of  $q$ ,  $q$ -de Rham cohomology in mixed characteristic is independent of choices, which was already known after base change to a period ring, via the comparisons of [BMS] between  $q$ -de Rham cohomology and their theory  $A\Omega^\bullet$ .

We expect that the dependence of these cohomology theories either on Adams operations at the residue characteristics (for de Rham) or on  $p$ -power roots of  $q$  (for variants of de Rham–Witt) is unavoidable, and that the conjectures of [Sch2] might thus be slightly

optimistic. Some of the strongest evidence for the conjectures is provided by the lifts of the Cartier isomorphisms, which rely on a choice of Frobenius. On the other hand, the comparison theorems of [BMS] can be seen as a manifestation of  $q$ -de Rham–Witt complexes; although they do not require a lift of Frobenius, they involve all  $p$ -power roots of  $q$ .

The essence of our construction of  $q$ -de Rham cohomology of  $A$  over  $R$  is to set  $q$  to be an element of rank 1 for the  $\Lambda$ -ring structure, and to look at flat  $\Lambda$ -rings  $B$  over  $R[q]$  equipped with morphisms  $A \rightarrow B/(q-1)$  of  $\Lambda$ -rings over  $R$ . If these seem unfamiliar, reassurance should be provided by the observation that  $(q-1)B$  carries  $q$ -analogues of divided power operations (Remark 1.4).

For the variants of de Rham–Witt cohomology in §3, the key to giving a characterisation independent of lifts of Frobenius is the factorisation of the tilting equivalence for perfectoid algebra via a category of  $\Lambda_p$ -rings.

I would like to thank Peter Scholze for many helpful comments, in particular about the possibility of a  $q$ -analogue of de Rham–Witt cohomology, and Michel Gros for spotting a missing hypothesis.

## CONTENTS

Introduction	1
1. Comparisons for $\Lambda$ -rings	2
1.1. The $\Lambda$ -ring $\mathbb{Z}[q]$	3
1.2. $q$ -cohomology of $\Lambda$ -rings	4
1.3. Completed $q$ -cohomology	6
2. Comparisons for $\Lambda_p$ -rings	8
2.1. $q$ -cohomology for $\Lambda_p$ -rings	8
2.2. Cartier isomorphisms in mixed characteristic	10
3. Functoriality via analogues of de Rham–Witt cohomology	11
3.1. Motivation	12
3.2. Almost isomorphisms	13
3.3. Perfectoid algebras	14
3.4. Functoriality of $q$ -de Rham cohomology	14
References	17

## 1. COMPARISONS FOR $\Lambda$ -RINGS

We will follow standard notational conventions for  $\Lambda$ -rings. These are commutative rings equipped with operations  $\lambda^i$  resembling alternating powers, in particular satisfying  $\lambda^k(a+b) = \sum_{i=0}^k \lambda^i(a)\lambda^{k-i}(b)$ , with  $\lambda^0(a) = 1$  and  $\lambda^1(a) = a$ . For background, see [Bor] and references therein. The  $\Lambda$ -rings we encounter are all torsion-free, in which case the  $\Lambda$ -ring structure is equivalent to giving ring endomorphisms  $\Psi^n$  for  $n \in \mathbb{Z}_{>0}$  with  $\Psi^{mn} = \Psi^m \circ \Psi^n$  and  $\Psi^p(x) \equiv x^p \pmod{p}$  for all primes  $p$ . If we write  $\lambda_t(f) := \sum_{i \geq 0} \lambda^i(f)t^i$  and  $\Psi_t(f) := \sum_{n \geq 1} \Psi^n(f)t^n$ , then the families of operations are related by the formula  $\Psi_t = -t \frac{d \log \lambda_t}{dt}$ .

We refer to elements  $x$  with  $\lambda^i(x) = 0$  for all  $i > 1$  (or equivalently  $\Psi^n(x) = x^n$  for all  $n$ ) as elements of rank 1.

### 1.1. The $\Lambda$ -ring $\mathbb{Z}[q]$ .

**Definition 1.1.** Define  $\mathbb{Z}[q]$  to be the  $\Lambda$ -ring with operations determined by setting  $q$  to be of rank 1.

We now consider the  $q$ -analogues  $[n]_q := \frac{q^n - 1}{q - 1} \in \mathbb{Z}[q]$  of the integers, with  $[n]_q! = [n]_q [n-1]_q \dots [1]_q$ , and  $\binom{n}{k}_q = \frac{[n]_q!}{[n-k]_q! [k]_q!}$ .

*Remark 1.2.* To see the importance of regarding  $\mathbb{Z}[q]$  as a  $\Lambda$ -ring observe that the binomial expressions

$$\lambda^k(n) = \binom{n}{k}, \quad \lambda^k(-n) = (-1)^k \binom{n+k-1}{k}$$

have as  $q$ -analogues the Gaussian binomial theorems

$$\lambda^k([n]_q) = q^{k(k-1)/2} \binom{n}{k}_q, \quad \lambda^k(-[n]_q) = (-1)^k \binom{n+k-1}{k}_q,$$

as well as Adams operations

$$\Psi^i([n]_q) = [n]_{q^i}.$$

For any torsion-free  $\Lambda$ -ring, localisation at a set of elements closed under the Adams operations always yields another  $\Lambda$ -ring, since  $\Psi^p(a^{-1}) - a^p = (\Psi^p(a)a^p)^{-1}(a^p - \Psi^p(a))$  is divisible by  $p$ .

**Lemma 1.3.** *For the  $\Lambda$ -ring structure on  $\mathbb{Z}[x, y]$  with  $x, y$  of rank 1, the elements*

$$\lambda^n\left(\frac{y-x}{q-1}\right) \in \mathbb{Z}[q, \{(q^n - 1)^{-1}\}_{n \geq 1}, x, y]$$

*are given by*

$$\begin{aligned} \lambda^k\left(\frac{y-x}{q-1}\right) &= \frac{(y-x)(y-qx) \dots (y-q^{k-1}x)}{(q-1)^k [k]_q!}, \\ &= \sum_{j=0}^k \frac{q^{j(j-1)/2} (-x)^j y^{k-j}}{[j]_q! [k-j]_q!}. \end{aligned}$$

*Proof.* The second expression comes from multiplying out the Gaussian binomial expansions. The easiest way to prove the first is to observe that  $\lambda^k(\frac{y-x}{q-1})$  must be a homogeneous polynomial of degree  $k$  in  $x, y$ , with coefficients in the integral domain  $\mathbb{Z}[q, \{(q^n - 1)^{-1}\}_{n \geq 1}]$ , and to note that

$$\lambda^k\left(\frac{q^n x - x}{q-1}\right) = \lambda^k([n]_q x) = q^{k(k-1)/2} \binom{n}{k}_q x^k.$$

Thus  $\lambda^k(\frac{y-x}{q-1})$  agrees with the homogeneous polynomial above for infinitely many values of  $\frac{y}{x}$ , so must be equal to it.  $\square$

*Remark 1.4.* Note that as  $q \rightarrow 1$ , Lemma 1.3 gives  $(q-1)^k \lambda^k(\frac{y-x}{q-1}) \rightarrow \frac{(x-y)^k}{k!}$ . Indeed, for any rank 1 element  $x$  in a  $\Lambda$ -ring we have  $\lambda_{(q-1)t}(\frac{x}{q-1}) = e_q(xt)$ , the  $q$ -exponential, with multiplicativity and universality then implying that  $\lambda_{(q-1)t}(\frac{a}{q-1})$  is a  $q$ -deformation of  $\exp(at)$  for all  $a$ . Thus  $(q-1)^k \lambda^k(\frac{a}{q-1})$  is a  $q$ -analogue of the  $k$ th divided power  $(a^k/k!)$ . An explicit expression comes recursively from the formula

$$[k]_q (q-1) \lambda^k\left(\frac{a}{q-1}\right) = \sum_{i>0} \lambda^i(a) \lambda^{k-i}\left(\frac{a}{q-1}\right).$$

**Lemma 1.5.** *For elements  $x, y$  of rank 1, the  $\Lambda$ -subring of  $\mathbb{Z}[q, \{(q^n - 1)^{-1}\}_{n \geq 1}, x, y]$  generated by  $q, x, y, \frac{y-x}{q-1}$  has basis  $\lambda^k(\frac{y-x}{q-1})$  as a  $\mathbb{Z}[q, x]$ -module.*

*Proof.* The  $\Lambda$ -subring clearly contains the  $\mathbb{Z}[q, x]$ -module  $M$  generated by the elements  $\lambda^k(\frac{y-x}{q-1})$ , which are also clearly linearly independent. Since  $\mathbb{Z}[x, q]$  is a  $\Lambda$ -ring, it suffices to show that  $M$  is closed under multiplication.

By Lemma 1.3, we know that

$$\lambda^i(\frac{y-x}{q-1})\lambda^j(\frac{y-q^i x}{q-1}) = \binom{i+j}{i}_q \lambda^{i+j}(\frac{y-x}{q-1}).$$

We can rewrite  $\frac{y-q^i x}{q-1} = \frac{y-x}{q-1} - [i]_q x$ , so  $\lambda^j(\frac{y-q^i x}{q-1}) - \lambda^j(\frac{y-x}{q-1})$  lies in the  $\mathbb{Z}[q, x]$ -module spanned by  $\lambda^m(\frac{y-x}{q-1})$  for  $m < j$ . By induction on  $j$ , it thus follows that

$$\lambda^i(\frac{y-x}{q-1})\lambda^j(\frac{y-q^i x}{q-1}) - \lambda^i(\frac{y-x}{q-1})\lambda^j(\frac{y-x}{q-1}) \in M,$$

so the binomial expression above implies  $\lambda^i(\frac{y-x}{q-1})\lambda^j(\frac{y-x}{q-1}) \in M$ .  $\square$

## 1.2. $q$ -cohomology of $\Lambda$ -rings.

**Definition 1.6.** Given a  $\Lambda$ -ring  $R$ , say that  $A$  is a  $\Lambda$ -ring over  $R$  if it is a  $\Lambda$ -ring equipped with a morphism  $R \rightarrow A$  of  $\Lambda$ -rings. We say that  $A$  is a flat  $\Lambda$ -ring over  $R$  if  $A$  is flat as a module over the commutative ring underlying  $R$ .

**Definition 1.7.** Given a morphism  $R \rightarrow A$  of  $\Lambda$ -rings, we define the category  $\text{Strat}_{A/R}^q$  to consist of flat  $\Lambda$ -rings  $B$  over  $R[q]$  equipped with a compatible morphism  $A \rightarrow B/(q-1)$ , such that the map  $A \rightarrow B/(q-1)$  admits a lift to  $B$ ; a choice of lift is not taken to be part of the data, so need not be preserved by morphisms.

More concisely,  $\text{Strat}_{A/R}^q$  is the Grothendieck construction of the functor

$$(\text{Spec } A)_{\text{strat}}^q: B \mapsto \text{Im}(\text{Hom}_{\Lambda, R}(A, B) \rightarrow \text{Hom}_{\Lambda, R}(A, B/(q-1)))$$

on the category  $f\Lambda(R[q])$  of flat  $\Lambda$ -rings over  $R[q]$ .

**Definition 1.8.** Given a flat morphism  $R \rightarrow A$  of  $\Lambda$ -rings, define  $\text{qDR}(A/R)$  to be the cochain complex of  $R[q]$ -modules given by taking the homotopy limit of the functor

$$\begin{aligned} \text{Strat}_{A/R}^q &\rightarrow \text{Ch}(R[q]) \\ B &\mapsto B. \end{aligned}$$

Equivalently, can we follow the approach of [Gro, Sim] towards the de Rham stack by regarding  $\text{qDR}(A/R)$  as the quasi-coherent cohomology complex of  $(\text{Spec } A)_{\text{strat}}^q$ . Writing  $\emptyset: f\Lambda(R[q]) \rightarrow \text{Mod}_{R[q]}$  for the forgetful functor to the category of  $R[q]$ -modules, and  $[f\Lambda(R[q]), \text{Set}]$  for the category of set-valued functors on  $f\Lambda(R[q])$ , we have

$$\text{qDR}(A/R) = \mathbf{R}\text{Hom}_{[f\Lambda(R[q]), \text{Set}]}((\text{Spec } A)_{\text{strat}}^q, \emptyset),$$

coming from the right-derived functor of the functor  $\text{Hom}_{[f\Lambda(R[q]), \text{Set}]}((\text{Spec } A)_{\text{strat}}^q, -)$  of natural transformations with source  $(\text{Spec } A)_{\text{strat}}^q$ .

**Definition 1.9.** Given a polynomial ring  $R[x]$ , recall from [Sch2] that the  $q$ -de Rham (or Aomoto–Jackson) cohomology  $q\text{-}\Omega_{R[x]/R}^\bullet$  is given by the complex

$$R[x][q] \xrightarrow{\nabla_q} R[x][q]dx, \quad \text{where} \quad \nabla_q(f) = \frac{f(qx) - f(x)}{x(q-1)}dx,$$

so  $\nabla_q(x^n) = [n]_q x^{n-1} dx$ .

Given a polynomial ring  $R[x_1, \dots, x_d]$ , the  $q$ -de Rham complex  $q\text{-}\Omega_{R[x_1, \dots, x_d]/R}^\bullet$  is then set to be

$$q\text{-}\Omega_{R[x_1]/R}^\bullet \otimes_{R[q]} q\text{-}\Omega_{R[x_2]/R}^\bullet \otimes_{R[q]} \cdots \otimes_{R[q]} q\text{-}\Omega_{R[x_d]/R}^\bullet,$$

so takes the form

$$R[x_1, \dots, x_d][q] \xrightarrow{\nabla_q} \Omega_{R[x_1, \dots, x_d]/R}^1[q] \xrightarrow{\nabla_q} \cdots \xrightarrow{\nabla_q} \Omega_{R[x_1, \dots, x_d]/R}^d[q].$$

**Proposition 1.10.** *If  $R$  is a  $\Lambda$ -ring and  $x$  of rank 1, then  $\text{qDR}(R[x]/R)$  can be calculated by a cosimplicial module  $U^\bullet$  given by setting  $U^n$  to be the  $\Lambda$ -subring*

$$U^n \subset R[q, \{(q^m - 1)^{-1}\}_{m \geq 1}, x_0, \dots, x_n]$$

*generated by  $q$  and the elements  $x_i$  and  $\frac{x_i - x_j}{q-1}$ .*

*Proof.* For  $X = \text{Spec } R[x]$ , the set-valued functor  $X_{\text{strat}}^q$  is not representable, but it can be resolved by the simplicial functor  $\tilde{X}_{\text{strat}}^q$  given by taking the Čech nerve of  $\text{Hom}_{\Lambda, R}(A, B) \rightarrow \text{Hom}_{\Lambda, R}(A, B/(q-1))$ , so

$$\begin{aligned} (\tilde{X}_{\text{strat}}^q)_n(B) &:= \overbrace{\text{Hom}_{\Lambda, R}(A, B) \times_{\text{Hom}_{\Lambda, R}(A, B/(q-1))} \cdots \times_{\text{Hom}_{\Lambda, R}(A, B/(q-1))} \text{Hom}_{\Lambda, R}(A, B)}^{n+1} \\ &= \text{Hom}_{\Lambda, R}(A, \overbrace{B \times_{B/(q-1)} \cdots \times_{B/(q-1)} B}^{n+1}). \end{aligned}$$

Observe that any element of  $(X_{\text{strat}}^q)_n(B)$  gives rise to a morphism  $f: R[q, x_0, \dots, x_n] \rightarrow B$  of  $\Lambda$ -rings over  $R[q]$ , with the image of  $x_i - x_j$  divisible by  $(q-1)$ . Flatness of  $B$  then gives a unique element  $f(x_i - x_j)/(q-1) \in B$ , so we have a map  $f$  to  $B$  from the free  $\Lambda$ -ring  $L$  over  $R[q, x_0, \dots, x_n]$  generated by elements  $z_{ij}$  with  $(q-1)z_{ij} = x_i - x_j$ .

Since  $B$  is flat, it embeds in  $B[\{(q^m - 1)^{-1}\}_{m \geq 1}]$  (the only hypothesis we really need) implying that the image of  $f$  factors through the image  $U^n$  of  $L$  in  $R[q, \{(q^m - 1)^{-1}\}_{m \geq 1}, x_0, \dots, x_n]$ . To see that  $(X_{\text{strat}}^q)_n$  is represented by  $U^n$ , we only now need to check that  $U^n$  is itself flat over  $R[q]$ , which follows because the argument of Lemma 1.5 gives a basis

$$x_0^{r_0} \lambda^{r_1} \left( \frac{x_1 - x_0}{q-1} \right) \cdots \lambda^{r_n} \left( \frac{x_n - x_{n-1}}{q-1} \right)$$

for  $U^n$  over  $R[q]$ . We therefore have  $\text{qDR}(R[x]/R) \simeq U^\bullet$ .  $\square$

**Theorem 1.11.** *If  $R$  is a  $\Lambda$ -ring, and the polynomial ring  $R[x_1, \dots, x_d]$  is given the  $\Lambda$ -ring structure for which the elements  $x_i$  are of rank 1, then there are  $R[q]$ -linear zigzags of quasi-isomorphisms*

$$\begin{aligned} \text{qDR}(R[x_1, \dots, x_n]/R) &\simeq (\Omega_{R[x_1, \dots, x_n]/R}^*[q], (q-1)\nabla_q) \\ \mathbf{L}\eta_{(q-1)} \text{qDR}(R[x_1, \dots, x_n]/R) &\simeq q\text{-}\Omega_{R[x_1, \dots, x_n]/R}^\bullet, \end{aligned}$$

where  $\mathbf{L}\eta_{(q-1)}$  denotes derived décalage with respect to the  $(q-1)$ -adic filtration.

*Proof.* It suffices to prove the first statement, the second following immediately by décalage. Since  $(\text{Spec } A \otimes_R A')_{\text{strat}}^q(B) = (\text{Spec } A)_{\text{strat}}^q(B) \times (\text{Spec } A')_{\text{strat}}^q(B)$ , and co-product of flat  $\Lambda$ -rings over  $R[q]$  is given by  $\otimes_{R[q]}$ , we have  $\text{qDR}((A \otimes_R A')/R) \simeq \text{qDR}(A/R) \otimes_{R[q]}^{\mathbf{L}} \text{qDR}(A'/R)$ , so we may reduce to the case  $A = R[x]$ .

Proposition 1.10 gives  $\mathrm{qDR}(R[x]/R) \simeq U^\bullet$ , and in order to compare this with  $q$ -de Rham cohomology, we now consider the cochain complexes  $\tilde{\Omega}^\bullet(U^n)$  given by

$$U^n \xrightarrow{(q-1)\nabla_q} \bigoplus_i U^n dx_i \xrightarrow{(q-1)\nabla_q} \bigoplus_{i < j} U^n dx_i \wedge dx_j \xrightarrow{(q-1)\nabla_q} \dots$$

In order to see that this differential is well-defined, observe that

$$\begin{aligned} (q-1)\nabla_{q,y} \lambda^k\left(\frac{y-x}{q-1}\right) &= y^{-1}(\lambda^k\left(\frac{qy-x}{q-1}\right) - \lambda^k\left(\frac{y-x}{q-1}\right))dy \\ &= y^{-1}(\lambda^k(y + \frac{y-x}{q-1}) - \lambda^k(\frac{y-x}{q-1}))dy \\ &= \lambda^{k-1}\left(\frac{y-x}{q-1}\right)dy, \end{aligned}$$

and similarly

$$(q-1)\nabla_{q,x} \lambda^k\left(\frac{y-x}{q-1}\right) = \sum_{i \geq 1} (-1)^i x^{i-1} \lambda^{k-i}\left(\frac{y-x}{q-1}\right) dx.$$

The first calculation also shows that the inclusion  $\tilde{\Omega}^\bullet(U^{n-1}) \hookrightarrow \tilde{\Omega}^\bullet(U^n)$  is a quasi-isomorphism, since

$$(q-1)\nabla_{q,x_n}(f(x_0, \dots, x_{n-1})\lambda^k(\frac{x_n-x_{n-1}}{q-1})) = f(x_0, \dots, x_{n-1})\lambda^{k-1}(\frac{x_n-x_{n-1}}{q-1})dx_n.$$

By induction on  $n$  we deduce that the inclusion  $\tilde{\Omega}^\bullet(U^0) \hookrightarrow \tilde{\Omega}^\bullet(U^n)$ , and hence the retraction of it given by the diagonal, is a quasi-isomorphism.

Now the complexes  $\tilde{\Omega}^i(U^\bullet)$  are all acyclic for  $i > 0$ , consisting of cosimplicial tensor products of  $U^\bullet$  with cosimplicial symmetric powers of the acyclic complex given by  $\mathbb{Z}dx_0 \oplus \dots \oplus \mathbb{Z}dx_n$  in level  $n$ . We therefore have quasi-isomorphisms

$$U^\bullet \leftarrow \mathrm{Tot} \tilde{\Omega}^\bullet(U^\bullet) \rightarrow \tilde{\Omega}^\bullet(U^0)$$

of flat cochain complexes over  $R[q]$ , so

$$\mathrm{qDR}(R[x]/R) \simeq \tilde{\Omega}^\bullet(R[x]),$$

and we just observe that  $\eta_{(q-1)}\tilde{\Omega}^\bullet(R[x]) = (\Omega_{R[x]/R}^*[q], (q-1)\nabla_q)$ .  $\square$

*Remark 1.12.* Note that Theorem 1.11 implies that  $q\text{-}\Omega_{R[x_1, \dots, x_n]/R}^\bullet$  naturally underlies the décalage of a cosimplicial  $\Lambda$ -ring over  $R[q]$ . Even the underlying cosimplicial commutative ring structure carries more information than an  $E_\infty$ -structure when  $\mathbb{Q} \not\subseteq R$ .

*Remark 1.13.* The complex  $(\Omega_{R[x_1, \dots, x_n]/R}^*[q], (q-1)\nabla_q)$  is a more fundamental object than its décalage  $q\text{-}\Omega_{R[x_1, \dots, x_n]/R}^\bullet$ , since it has a vestigial memory of the Hodge filtration.

There might be a natural formulation of the theorem not involving décalage, in terms of a  $q$ -analogue of the crystalline site for a  $\Lambda$ -ring  $A$  over  $R$ , regarded as an  $R[q]$ -algebra via  $R = R[q]/(q-1)$ . Following Remark 1.4, this would involve extensions  $B \rightarrow A$  of  $\Lambda$ -rings over  $R[q]$  equipped with  $q$ -analogues of divided power operations on the augmentation ideals  $I$ , looking like  $x \mapsto (q-1)^k \lambda^k(\frac{x}{q-1})$ .

### 1.3. Completed $q$ -cohomology.

**Definition 1.14.** Given a morphism  $R \rightarrow A$  of  $\Lambda$ -rings, we define the category  $\hat{\mathrm{Strat}}_{A/R}^q \subset \mathrm{Strat}_{A/R}^q$  to consist of those objects which are  $(q-1)$ -adically complete.

**Definition 1.15.** Given a flat morphism  $R \rightarrow A$  of  $\Lambda$ -rings, define  $q\widehat{\mathrm{DR}}(A/R)$  to be the cochain complex of  $R[[q-1]]$ -modules given by taking the homotopy limit of the functor

$$\begin{aligned} \widehat{\mathrm{Strat}}_{A/R}^q &\rightarrow \mathrm{Ch}(R[[q]]) \\ B &\mapsto B. \end{aligned}$$

The following is immediate:

**Lemma 1.16.** *Given a flat morphism  $R \rightarrow A$  of  $\Lambda$ -rings, the complex  $q\widehat{\mathrm{DR}}(A/R)$  is the derived  $(q-1)$ -adic completion of  $q\mathrm{DR}(A/R)$ .*

**Definition 1.17.** As in [Sch2, §3], given a formally étale map  $\square: R[x_1, \dots, x_d] \rightarrow A$ , define  $\widehat{q\text{-}\Omega}_{A/R, \square}^\bullet$  to be the complex

$$A[[q-1]] \xrightarrow{\nabla_q} \Omega_{A/R}^1[[q-1]] \xrightarrow{\nabla_q} \dots \xrightarrow{\nabla_q} \Omega_{A/R}^d[[q-1]],$$

where  $\nabla_q$  is defined as follows. First note that the ring endomorphisms  $\gamma_i$  of  $R[x_1, \dots, x_d][[q-1]]$  given by  $\gamma_i(x_j) = q^{\delta_{ij}}x_j$  extend uniquely to endomorphisms of  $A[[q-1]]$  which are the identity modulo  $(q-1)$ , then set

$$\nabla_q(f) := \sum_i \frac{\gamma_i(f) - f}{(q-1)x_i} dx_i.$$

Note that  $\widehat{q\text{-}\Omega}_{R[x_1, \dots, x_d]/R}^\bullet$  is just the  $(q-1)$ -adic completion of  $q\text{-}\Omega_{R[x_1, \dots, x_d]/R}^\bullet$ .

**Proposition 1.18.** *If  $R$  is a flat  $\Lambda$ -ring over  $\mathbb{Z}$  and  $\square: R[x_1, \dots, x_d] \rightarrow A$  is a formally étale map of  $\Lambda$ -rings, the elements  $x_i$  having rank 1, then there are zigzags of  $R[[q]]$ -linear quasi-isomorphisms*

$$q\widehat{\mathrm{DR}}(A/R) \simeq (\Omega_{A/R}^*[[q-1]], (q-1)\nabla_q), \quad \mathbf{L}\eta_{(q-1)}q\widehat{\mathrm{DR}}(A/R) \simeq \widehat{q\text{-}\Omega}_{A/R, \square}^\bullet.$$

The induced quasi-isomorphisms

$$q\widehat{\mathrm{DR}}(A/R) \otimes_{R[[q-1]]}^{\mathbf{L}} R \simeq (\Omega_{A/R}^*, 0), \quad (\mathbf{L}\eta_{(q-1)}q\widehat{\mathrm{DR}}(A/R)) \otimes_{R[[q-1]]}^{\mathbf{L}} R \simeq \Omega_{A/R}^\bullet$$

are independent of the choice of framing.

*Proof.* This is much the same as the proof of Theorem 1.11. The complex  $q\widehat{\mathrm{DR}}(A/R)$  can be realised as a cosimplicial  $\Lambda$ -ring  $U$ , with  $U^n$  the  $(q-1)$ -adically complete  $\Lambda$ -subring of  $A^{\otimes_R(n+1)}[[q-1]][\{(q^m-1)^{-1}\}_{m \geq 1}]$  generated by

$$A^{\otimes_R(n+1)}[[q-1]] \quad \text{and} \quad (q-1)^{-1} \ker(A^{\otimes_R(n+1)} \rightarrow A)[[q-1]].$$

Uniqueness of lifts with respect to the formally étale framing ensures that the endomorphisms  $\gamma_i$  commute with the Adams operations, so are  $\Lambda$ -ring endomorphisms of  $R$ . Since the formal completion of  $A \otimes_R A \rightarrow A$  is just the  $\Lambda$ -ring

$$A[(x_1 - y_1), (x_2 - y_2), \dots, (x_d - y_d)],$$

the calculations of Theorem 1.11 now adapt to give quasi-isomorphisms

$$(\Omega_{A/R}^*[[q]], (q-1)\nabla_q) \leftarrow \mathrm{Tot} \tilde{\Omega}^\bullet(U^\bullet) \rightarrow U^\bullet,$$

where  $\tilde{\Omega}^\bullet(U^n)$  is the  $(q-1)$ -adic completion of  $(U^n \otimes_{A^{\otimes(n+1)}} (\Omega_{A/R}^*)^{\otimes(n+1)}, (q-1)\nabla_q)$ . Reduction of this or its décalage modulo  $(q-1)$  replaces  $\nabla_q$  with  $d$ , removing the dependence on co-ordinates.  $\square$



*Remark 1.19.* As in [Sch2, Definition 7.3], there is a notion of  $q$ -connections on projective  $A[[q-1]]$ -modules  $M$ . Adapting the ideas of Proposition 1.18, these will be equivalent to projective modules over  $X_{\text{strat}}^q$ , so flat Cartesian  $\widehat{\text{qDR}}(A/R)$ -modules  $N$  with  $N \otimes_{\widehat{\text{qDR}}(A/R)} A[[q-1]] = M$ , together with a condition that the  $(\widehat{\text{qDR}}(A/R)/(q-1))$ -module  $N/(q-1)$  is just given by pullback of the  $A$ -module  $M/(q-1)$ .

Via Lemma 1.5, these data are equivalent to specifying an operator  $\partial^1: M \rightarrow \bigoplus_{k_1, \dots, k_d} M \lambda^{k_1}(\frac{y_1-x_1}{q-1}) \dots \lambda^{k_d}(\frac{y_d-x_d}{q-1})$  satisfying a cocycle condition and congruent to the identity modulo  $(q-1)$ . Such operators then arise from  $q$ -connections  $(\nabla_{1,q}, \dots, \nabla_{d,q})$  as  $q$ -Taylor series

$$\partial^1(f) := \sum_{k_1, \dots, k_d} (q-1)^{\sum k_i} (\nabla_{1,q}^{k_1} \dots \nabla_{d,q}^{k_d})(f) \lambda^{k_1}(\frac{y_1-x_1}{q-1}) \dots \lambda^{k_d}(\frac{y_d-x_d}{q-1}).$$

## 2. COMPARISONS FOR $\Lambda_P$ -RINGS

Since very few étale maps  $R[x_1, \dots, x_d] \rightarrow A$  give rise to  $\Lambda$ -ring structures on  $A$ , Proposition 1.18 is fairly limited in its scope for applications. We now show how the construction of  $\widehat{\text{qDR}}$  and the comparison quasi-isomorphism survive when we weaken the  $\Lambda$ -ring structure by discarding Adams operations at invertible primes.

**2.1.  $q$ -cohomology for  $\Lambda_P$ -rings.** Our earlier constructions for  $\Lambda$ -rings all carry over to  $\Lambda_P$ -rings, as follows.

**Definition 2.1.** Given a set  $P$  of primes, we define a  $\Lambda_P$ -ring  $A$  to be a  $\Lambda_{\mathbb{Z},P}$ -ring in the sense of [Bor]. This means that it is a coalgebra in commutative rings for the comonad given by the functor  $W^{(P)}$  of  $P$ -typical Witt vectors. When a commutative ring  $A$  is flat over  $\mathbb{Z}$ , giving a  $\Lambda_P$ -ring structure on  $A$  is equivalent to giving commuting Adams operations  $\Psi^p$  for all  $p \in P$ , with  $\Psi^p(a) \equiv a^p \pmod{p}$  for all  $a$ .

Thus when  $P$  is the set of all primes, a  $\Lambda_P$ -ring is just a  $\Lambda$ -ring; a  $\Lambda_\emptyset$ -ring is just a commutative ring; for a single prime  $p$ , we write  $\Lambda_p := \Lambda_{\{p\}}$ , and note that a  $\Lambda_p$ -ring is a  $\delta$ -ring in the sense of [Joy].

**Definition 2.2.** Given a  $\Lambda_P$ -ring  $R$ , say that  $A$  is a  $\Lambda_P$ -ring over  $R$  if it is a  $\Lambda_P$ -ring equipped with a morphism  $R \rightarrow A$  of  $\Lambda_P$ -rings. We say that  $A$  is a flat  $\Lambda_P$ -ring over  $R$  if  $A$  is flat as a module over the commutative ring underlying  $R$ .

**Definition 2.3.** Given a morphism  $R \rightarrow A$  of  $\Lambda_P$ -rings, we define the category  $\text{Strat}_{A/R}^{q,P}$  to consist of flat  $\Lambda_P$ -rings  $B$  over  $R[q]$  equipped with a compatible morphism  $A \rightarrow B/(q-1)$ , such that the map  $A \rightarrow B/(q-1)$  admits a lift to  $B$ . We define the category  $\widehat{\text{Strat}}_{A/R}^{q,P} \subset \text{Strat}_{A/R}^q$  to consist of those objects which are  $(q-1)$ -adically complete.

More concisely,  $\text{Strat}_{A/R}^{q,P}$  (resp.  $\widehat{\text{Strat}}_{A/R}^{q,P}$ ) is the Grothendieck construction of the functor

$$(\text{Spec } A)_{\text{strat}}^{q,P}: B \mapsto \text{Im}(\text{Hom}_{\Lambda_P, R}(A, B) \rightarrow \text{Hom}_{\Lambda_P, R}(A, B/(q-1)))$$

of the category of flat  $\Lambda_P$ -rings (resp.  $(q-1)$ -adically complete flat  $\Lambda_P$ -rings) over  $R[q]$ .

**Definition 2.4.** Given a flat morphism  $R \rightarrow A$  of  $\Lambda_P$ -rings, define  $\mathrm{qDR}_P(A/R)$  to be the cochain complex of  $R[q]$ -modules given by taking the homotopy limit of the functor

$$\begin{aligned} \mathrm{Strat}_{A/R}^{q,P} &\rightarrow \mathrm{Ch}(R[q]) \\ B &\mapsto B. \end{aligned}$$

Define  $\hat{\mathrm{qDR}}_P(A/R)$  to be the cochain complex of  $R[[q-1]]$ -modules given by the corresponding homotopy limit over  $\hat{\mathrm{Strat}}_{A/R}^{q,P}$ .

Thus when  $P$  is the set of all primes, we have  $\mathrm{qDR}_P(A/R) = \mathrm{qDR}(A/R)$ . At the other extreme, for  $A$  smooth,  $\hat{\mathrm{qDR}}_\emptyset(A/R)$  is the Rees construction of the Hodge filtration on the infinitesimal cohomology complex of  $A$  over  $R$ , with formal variable  $(q-1)$ . In more detail, there is a decreasing filtration  $F$  of  $\mathcal{O}_{\mathrm{inf}}$  given by powers of the augmentation ideal  $\mathcal{O}_{\mathrm{inf}} \rightarrow \mathcal{O}_{\mathrm{Zar}}$ , and  $\hat{\mathrm{qDR}}_\emptyset(A/R) \simeq \bigoplus_{\nu \in \mathbb{Z}} (q-1)^{-\nu} \mathbf{R}\Gamma(\mathrm{Spec} A, F^\nu \mathcal{O}_{\mathrm{inf}})(q-1)^{-\nu}$ .

**Lemma 2.5.** *For a set  $P$  of primes, the forgetful functor from  $\Lambda_P$ -rings to  $\Lambda$ -rings has a right adjoint  $W^{(\notin P)}$ . There is a canonical ghost component morphism*

$$W^{(\notin P)}(B) \rightarrow \prod_{\substack{n \in \mathbb{N}: \\ (n,p)=1 \ \forall p \in P}} B,$$

which is an isomorphism when  $P$  contains all the residue characteristics of  $B$ .

*Proof.* Existence of a right adjoint follows from the comonadic definitions of  $\Lambda$ -rings and  $\Lambda_P$ -rings. The ghost component morphism is given by taking the Adams operations  $\Psi^n$  given by the  $\Lambda$ -ring structure on  $W^{(\notin P)}(B)$ , followed by projection to  $B$ . When  $P$  contains all the residue characteristics of  $B$ , a  $\Lambda$ -ring structure is the same as a  $\Lambda_P$ -ring structure with compatible commuting Adams operations for all primes not in  $P$ , leading to the description above.  $\square$

Note that the big Witt vector functor  $W$  on commutative rings thus factorises as  $W = W^{(\notin P)} \circ W^{(P)}$ , for  $W^{(P)}$  the  $P$ -typical Witt vectors.

**Proposition 2.6.** *Given a morphism  $R \rightarrow A$  of  $\Lambda$ -rings, and a set  $P$  of primes, there are natural maps*

$$\mathrm{qDR}_P(A/R) \rightarrow \mathrm{qDR}(A/R), \quad \hat{\mathrm{qDR}}_P(A/R) \rightarrow \hat{\mathrm{qDR}}(A/R),$$

and the latter map is a quasi-isomorphism when  $P$  contains all the residue characteristics of  $A$ .

*Proof.* We have functors

$$\begin{aligned} (\mathrm{Spec} A)_{\mathrm{strat}}^q \circ W^{(\notin P)}: B &\mapsto \mathrm{Im}(\mathrm{Hom}_{\Lambda, R}(A, W^{(\notin P)}B) \rightarrow \mathrm{Hom}_{\Lambda, R}(A, (W^{(\notin P)}B)/(q-1))) \\ (\mathrm{Spec} A)_{\mathrm{strat}}^{q,P}: B &\mapsto \mathrm{Im}(\mathrm{Hom}_{\Lambda_P, R}(A, B) \rightarrow \mathrm{Hom}_{\Lambda_P, R}(A, B/(q-1))) \end{aligned}$$

on the category of flat  $\Lambda_P$ -rings over  $R[q]$ . There is an obvious map

$$(W^{(\notin P)}B)/(q-1) \rightarrow W^{(\notin P)}(B/(q-1)),$$

and hence a natural transformation  $(\mathrm{Spec} A)_{\mathrm{strat}}^q \circ W^{(\notin P)} \rightarrow (\mathrm{Spec} A)_{\mathrm{strat}}^{q,P}$ , which induces the morphism  $\mathrm{qDR}_P(A/R) \rightarrow \mathrm{qDR}(A/R)$  on cohomology.

When  $P$  contains all the residue characteristics of  $A$ , the map  $(W^{(\notin P)}B)/(q-1) \rightarrow W^{(\notin P)}(B/(q-1))$  is just

$$\prod_{\substack{n \in \mathbb{N}: \\ (n,p)=1 \forall p \in P}} B/(q^n - 1) \rightarrow \prod_{\substack{n \in \mathbb{N}: \\ (n,p)=1 \forall p \in P}} B/(q-1),$$

since the morphism  $R[q] \rightarrow W^{(\notin P)}B$  is given by Adams operations, with  $\Psi^n(q-1) = q^n - 1$ .

We have  $(q^n - 1) = (q-1)[n]_q$ , and  $[n]_q$  is a unit in  $\mathbb{Z}[\frac{1}{n}][[q-1]]$ , hence a unit in  $B$  when  $n$  is coprime to the residue characteristics. Thus the map  $(W^{(\notin P)}B)/(q-1) \rightarrow W^{(\notin P)}(B/(q-1))$  gives an isomorphism whenever  $B$  is  $(q-1)$ -adically complete and admits a map from  $A$ , so the transformation  $(\text{Spec } A)_{\text{strat}}^q \circ W^{(\notin P)} \rightarrow (\text{Spec } A)_{\text{strat}}^{q,P}$  is a natural isomorphism on the category of flat  $(q-1)$ -adically complete  $\Lambda_P$ -rings over  $R[q]$ , and so  $\text{q}\widehat{\text{DR}}_P(A/R) \xrightarrow{\sim} \widehat{\text{qDR}}(A/R)$ .  $\square$

Over  $\mathbb{Z}[\frac{1}{P}]$ , every  $\Lambda_P$ -ring can be canonically made into a  $\Lambda$ -ring, by setting all the additional Adams operations to be the identity. However, this observation is of limited use in establishing functoriality of  $q$ -de Rham cohomology, because the resulting  $\Lambda$ -ring structure will not satisfy the conditions of Proposition 1.18. We now give a more general result which does allow for meaningful comparisons.

**Theorem 2.7.** *If  $R$  is a flat  $\Lambda_P$ -ring over  $\mathbb{Z}$  and  $\square: R[x_1, \dots, x_d] \rightarrow A$  is a formally étale map of  $\Lambda_P$ -rings, the elements  $x_i$  having rank 1, then there is a zigzag of  $R[[q]]$ -linear quasi-isomorphisms*

$$\text{L}\eta_{(q-1)}\widehat{\text{qDR}}_P(A/R) \simeq q\text{-}\widehat{\Omega}_{A/R, \square}^\bullet$$

whenever  $P$  contains all the residue characteristics of  $A$ .

*Proof.* The key observation to make is that formally étale maps have a unique lifting property with respect to nilpotent extensions of flat  $\Lambda_P$ -rings, because the Adams operations must also lift uniquely. In particular, this means that the operations  $\gamma_i$  featuring in the definition of  $q$ -de Rham cohomology are necessarily endomorphisms of  $A$  as a  $\Lambda_P$ -ring.

Similarly to Proposition 1.18,  $\widehat{\text{qDR}}_P(A/R)$  is calculated using a cosimplicial  $\Lambda_P$ -ring given in level  $n$  by the  $(q-1)$ -adic completion  $\hat{U}_{P,A}^\bullet$  of the  $\Lambda_P$ -ring over  $R[q]$  generated by  $A^{\otimes_R(n+1)}[q]$  and  $(q-1)^{-1} \ker(A^{\otimes_R(n+1)} \rightarrow A)[q]$ . The observation above shows that  $\hat{U}_{P,A}^n \cong \hat{U}_{P,R[x_1, \dots, x_d]}^n \hat{\otimes}_{R[x_1, \dots, x_d]} A$ , changing base along  $\square$  applied to the first factor.

As in Proposition 2.6,  $\hat{U}_{P,R[x_1, \dots, x_d]}^\bullet$  is just the  $(q-1)$ -adic completion of the complex  $U^\bullet$  from Proposition 1.10. Further application of the key observation above then allows us to adapt the constructions of Theorem 1.11, giving the desired quasi-isomorphisms.  $\square$

**2.2. Cartier isomorphisms in mixed characteristic.** The only setting in which Theorem 2.7 leads to results close to the conjectures of [Sch2] is when  $R = W^{(p)}(k)$ , the  $p$ -typical Witt vectors of a field of characteristic  $p$ , and  $A = \varprojlim_n A_n$  is a formal deformation of a smooth  $k$ -algebra  $A_0$ . Then any formally étale morphism  $W^{(p)}(k)[x_1, \dots, x_d] \rightarrow A$  gives rise to a unique compatible lift  $\Psi$  of absolute Frobenius on  $A$  with  $\Psi(x_i) = x_i^p$ , so gives  $A$  the structure of a topological  $\Lambda_p$ -ring. The framing still affects the choice of  $\Lambda_p$ -ring structure, but at least such a structure is

guaranteed to exist, giving rise to a complex  $\mathrm{qDR}_P(A/R)^{\wedge p} := \varprojlim_n \mathrm{qDR}_p(A/R) \otimes_R R_n$  depending only on the choice of  $\Psi$ , where  $R_n = W_n^{(p)}(k)$ .

Our constructions now allow us to globalise the quasi-isomorphism

$$(\widehat{q\text{-}\Omega}_{A/R, \square}^\bullet)^{\wedge p} / [p]_q \simeq (\Omega_{A/R}^*)^{\wedge p} \llbracket q-1 \rrbracket / [p]_q$$

of [Sch2, Proposition 3.4], where  $\Omega_{A/R}^*$  denotes the complex  $A \xrightarrow{0} \Omega_{A/R}^1 \xrightarrow{0} \Omega_{A/R}^2 \xrightarrow{0} \dots$

**Proposition 2.8.** *Take a smooth formal scheme  $\mathfrak{X}$  over  $R = W^{(p)}(k)$  equipped with a lift  $\Psi$  of Frobenius which étale locally admits co-ordinates  $\{x_i\}_i$  as above with  $\Psi(x_i) = x_i^p$ . Then there is a global quasi-isomorphism*

$$C_q^{-1}: (\Omega_{\mathfrak{X}/R}^*)^{\wedge p} \llbracket q-1 \rrbracket / [p]_q \rightarrow (\mathbf{L}\eta_{(q-1)} \mathrm{q}\hat{\mathrm{DR}}_p(\mathcal{O}_{\mathfrak{X}}/R))^{\wedge p} / [p]_q$$

in the derived category of étale sheaves on  $\mathfrak{X}$ .

*Proof.* Functoriality of the construction  $\mathrm{qDR}_p$  for rings with Frobenius lifts gives us a sheaf  $\mathrm{q}\hat{\mathrm{DR}}_p(\mathcal{O}_{\mathfrak{X}}/R)^{\wedge p}$  on  $\mathfrak{X}$ . We then have maps

$$\begin{aligned} \Psi^p: \mathrm{qDR}_p(\mathcal{O}_{\mathfrak{X}}/R)^{\wedge p} &\rightarrow \mathrm{qDR}_p(\mathcal{O}_{\mathfrak{X}}/R)^{\wedge p} \\ \mathrm{qDR}_p(\mathcal{O}_{\mathfrak{X}}/R)^{\wedge p} / (q-1) &\rightarrow \mathrm{qDR}_p(\mathcal{O}_{\mathfrak{X}}/R)^{\wedge p} / (q^p-1), \end{aligned}$$

and thus, denoting good truncation by  $\tau$ ,

$$(q-1)^i \Psi^p: \tau^{\leq i}(\mathrm{qDR}_p(\mathcal{O}_{\mathfrak{X}}/R)^{\wedge p} / (q-1)) \rightarrow (\mathbf{L}\eta_{(q-1)} \mathrm{q}\hat{\mathrm{DR}}_p(\mathcal{O}_{\mathfrak{X}}/R)^{\wedge p}) / [p]_q;$$

the left-hand side is quasi-isomorphic to  $\bigoplus_{j \leq i} (\Omega_{\mathcal{O}_{\mathfrak{X}}/R}^j)^{\wedge p} [-j]$  by Proposition 1.18.

Extending the construction  $R[q]$ -linearly and restricting to top summands therefore gives us the global map  $C_q^{-1}$ . For a local choice of framing, the map  $\Psi^p$  necessarily corresponds via Theorem 2.7 to the chain map  $adx^I \mapsto \Psi^p(a)x^{I(p-1)}dx^I$  on the complex  $(\Omega_{A/R}^* \llbracket q-1 \rrbracket, (q-1)\nabla_q)$ . This gives equivalences

$$(q-1)^i \Psi^p \simeq \sum_{j \leq i} (q-1)^{i-j} (\tilde{C}^{-1})^j$$

for Scholze's locally defined lifts  $(\tilde{C}^{-1})^j: (\Omega_{A/R}^j)^{\wedge p} [-j] \rightarrow (\widehat{q\text{-}\Omega}_{A/R, \square}^\bullet)^{\wedge p} / [p]_q$  of the Cartier quasi-isomorphism. The local calculation of [Sch2, Proposition 3.4] then ensures that  $C_q^{-1}$  is a quasi-isomorphism.  $\square$

### 3. FUNCTORIALITY VIA ANALOGUES OF DE RHAM–WITT COHOMOLOGY

In order to obtain a cohomology theory for smooth commutative rings rather than for  $\Lambda_p$ -rings, we now consider  $q$ -analogues of de Rham–Witt cohomology. Our starting point is to observe that if we allow roots of  $q$ , we can extend the Jackson differential to fractional powers of  $x$  by the formula

$$\nabla_q(x^{m/n}) = \frac{q^{m/n} - 1}{q - 1} x^{m/n} d \log x,$$

so terms such as  $[n]_{q^{1/n}} x^{m/n}$  have integral derivative, where  $[n]_{q^{1/n}} = \frac{q-1}{q^{1/n}-1}$ .

### 3.1. Motivation.

**Definition 3.1.** Given a  $\Lambda_P$ -ring  $B$ , define  $\Psi^{1/P^\infty} B$  to be the smallest  $\Lambda_P$ -ring containing  $B$  on which the Adams operations are automorphisms.

In the case  $P = \{p\}$ , the  $\Lambda_p$ -ring  $\Psi^{1/p^\infty} B$  is thus the colimit of the diagram

$$B \xrightarrow{\Psi^p} B \xrightarrow{\Psi^p} B \xrightarrow{\Psi^p} \dots$$

The proof of Theorem 2.7 allows us to replace  $\widehat{\mathrm{qDR}}_p(A/R)$  with the complex  $(\Omega_{A/R}^*[q-1], (q-1)\nabla_q)$ ; under this quasi-isomorphism, the Adams operations on  $A$  extend to  $\Omega_{A/R}^*[q-1]$  by setting  $\Psi^n(dx_i) := x_i^{n-1}dx_i$ . As an immediate consequence we have:

**Lemma 3.2.** *If  $R$  is a flat  $\Lambda_p$ -ring over  $\mathbb{Z}$  with  $\Psi^{1/p^\infty} R = R$  and residue characteristic  $p$ , then  $\Psi^{1/p^\infty} \mathrm{qDR}_p(R[x]/R) \simeq (\Omega_{R[x^{1/p^\infty}]/R}^*[q^{1/p^\infty}], (q-1)\nabla_q)$ , so the décalage  $\mathbf{L}\eta_{(q-1)} \Psi^{1/p^\infty} \widehat{\mathrm{qDR}}_p(R[x]/R)$  is quasi-isomorphic to the  $(q-1)$ -adic completion of the complex*

$$\begin{aligned} \{a \in R[x^{1/p^\infty}, q^{1/p^\infty}] : \nabla_q a \in R[x^{1/p^\infty}, q^{1/p^\infty}] d \log x\} &\xrightarrow{\nabla_q} \\ \{b d \log x \in R[x^{1/p^\infty}, q^{1/p^\infty}] d \log x : b(0, q) = 0\}. \end{aligned}$$

Thus in level 0 (resp. level 1),  $\mathbf{L}\eta_{(q-1)} \Psi^{1/p^\infty} \widehat{\mathrm{qDR}}_p(R[x]/R)$  is spanned by elements of the form  $[p^n]_{q^{1/p^n}} x^{m/p^n}$  (resp.  $x^{m/p^n} d \log x$ ), so setting  $q^{1/p^\infty} = 1$  gives a complex whose  $p$ -adic completion is the  $p$ -typical de Rham–Witt complex.

**Lemma 3.3.** *Let  $R$  and  $A$  be flat  $p$ -adically complete  $\Lambda_p$ -algebras over  $\mathbb{Z}_p$ , with  $\Psi^{1/p^\infty} R = R$  and, for elements  $x_i$  of rank 1, a map  $\square: R[x_1, \dots, x_d]^{\wedge p} \rightarrow A$  of  $\Lambda_p$ -rings which is a flat  $p$ -adic deformation of an étale map. Then the map*

$$(R[q^{1/p^\infty}] \otimes_{R[q]} \mathbf{L}\eta_{(q-1)} \widehat{\mathrm{qDR}}_p(A/R))^{\wedge p} \rightarrow \mathbf{L}\eta_{(q-1)} (\Psi^{1/p^\infty} \widehat{\mathrm{qDR}}_p(A/R))^{\wedge p}$$

*is a quasi-isomorphism.*

*Proof.* The map  $\Psi^p: A \otimes_{R[x_1, \dots, x_d]} R[x_1^{1/p}, \dots, x_d^{1/p}] \rightarrow A$  becomes an isomorphism on  $p$ -adic completion, because  $\square$  is flat and we have an isomorphism modulo  $p$ . Thus

$$\Psi^{1/p^\infty} A \cong A[x_1^{1/p^\infty}, \dots, x_d^{1/p^\infty}]^{\wedge p} := (A \otimes_{R[x_1, \dots, x_d]} R[x_1^{1/p^\infty}, \dots, x_d^{1/p^\infty}])^{\wedge p}$$

Combined with the calculation of Lemma 3.2, this gives us a quasi-isomorphism between  $(\Psi^{1/p^\infty} \widehat{\mathrm{qDR}}_p(A/R))^{\wedge p}$  and the  $(p, q-1)$ -adic completion of

$$\bigoplus_{\alpha} A[[q-1]] x_1^{\alpha_1} \dots x_d^{\alpha_d} dx^I,$$

where  $\alpha \in p^{-\infty} \mathbb{Z}^d$  such that  $0 \leq \alpha_i < 1$  if  $i \notin I$  and  $-1 < \alpha_i \leq 0$  if  $i \in I$ .

We then observe that the contributions to the décalage  $\eta_{(q-1)}$  from terms with  $\alpha \neq 0$  must be acyclic, via a contracting homotopy defined by the restriction to  $\eta_{(q-1)}$  of the  $q$ -integration map

$$f x_1^{\alpha_1} \dots x_d^{\alpha_d} dx^I \mapsto f x_1^{\alpha_1} \dots x_d^{\alpha_d} \sum_{i \in I} \pm x_i [\alpha_i]_q^{-1} dx^{(I \setminus i)},$$

where  $[\frac{m}{p^n}]_q^{-1} = [m]_{q^{1/p^n}}^{-1} [p^n]_{q^{1/p^n}}$  for  $m$  coprime to  $p$ , noting that  $[m]_{q^{1/p^n}}$  is a unit in  $\mathbb{Z}[q^{1/p^\infty}]^{\wedge(p, q-1)}$ .  $\square$

*Remark 3.4.* The endomorphism given on  $\Psi^{1/P^\infty} \hat{\mathrm{qDR}}_P(A/R)$  by

$$a \mapsto \Psi^{1/n}([n]_q a) = [n]_{q^{1/n}} \Psi^{1/n} a$$

descends to an endomorphism of  $H^0(\Psi^{1/P^\infty} \hat{\mathrm{qDR}}_P(A/R)/(q-1))$ , which we may denote by  $V_n$  because it mimics Verschiebung in the sense that  $\Psi^n V_n = n \cdot \mathrm{id}$ .

For  $A$  smooth over  $\mathbb{Z}$ , we then have

$$\begin{aligned} H^0(\Psi^{1/P^\infty} \hat{\mathrm{qDR}}_P(A/\mathbb{Z})/(q-1))/(V_p : p \in P) &\cong A[q^{1/P^\infty}]/([p]_{q^{1/p}} : p \in P) \\ &\cong A[\zeta_{P^\infty}], \end{aligned}$$

for  $\zeta_n$  a primitive  $n$ th root of unity.

By adjunction, this gives an injective map

$$H^0(\Psi^{1/P^\infty} \hat{\mathrm{qDR}}_P(A/\mathbb{Z})/(q-1)) \hookrightarrow W^{(P)} A[\zeta_{P^\infty}]$$

of  $\Lambda_P$ -rings, which becomes an isomorphism on completing  $\Psi^{1/P^\infty} \hat{\mathrm{qDR}}(A/\mathbb{Z})$  with respect to the system  $\{([n]_{q^{1/n}})\}_{n \in P^\infty}$  of ideals, where we write  $P^\infty$  for the set of integers whose prime factors are all in  $P$ . This implies that the cokernel is annihilated by all elements of  $(q^{1/P^\infty} - 1)$ , so leads us to consider almost mathematics as in [GR].

**3.2. Almost isomorphisms.** Combined with Lemma 3.3, Remark 3.4 allows us to regard  $\mathbf{L}\eta_{(q-1)} \Psi^{1/p^\infty} \hat{\mathrm{qDR}}_p(A/\mathbb{Z}_p)^{\wedge_p}$  as being almost a  $q^{1/p^\infty}$ -analogue of  $p$ -typical de Rham–Witt cohomology. (From now on, we consider only the case  $P = \{p\}$ .)

The ideal  $(q^{1/p^\infty} - 1)^{\wedge_{(p,q-1)}} = \ker(\mathbb{Z}[q^{1/p^\infty}]^{\wedge_{(p,q-1)}} \rightarrow \mathbb{Z}_p)$  is equal to the  $p$ -adic completion of its square, since we may write it as the kernel  $W^{(p)}(\mathfrak{m})$  of  $W^{(p)}(\mathbb{F}_p[q^{1/p^\infty}]^{\wedge_{(q-1)}}) \rightarrow W^{(p)}(\mathbb{F}_p)$ , for the idempotent maximal ideal  $\mathfrak{m} = ((q-1)^{1/p^\infty})^{\wedge_{(q-1)}}$  in  $\mathbb{F}_p[q^{1/p^\infty}]^{\wedge_{(q-1)}}$ . If we set  $h^{1/p^n}$  to be the Teichmüller element

$$[q^{1/p^n} - 1] = \lim_{r \rightarrow \infty} (q^{1/p^{nr}} - 1)^{p^r} \in \mathbb{Z}[q^{1/p^\infty}]^{\wedge_{(p,q-1)}},$$

then  $W^{(p)}(\mathfrak{m}) = (h^{1/p^\infty})^{\wedge_{(p,h)}}$ . Although  $W^{(p)}(\mathfrak{m})/p^n$  is not maximal in  $\mathbb{Z}[h^{1/p^\infty}]^{\wedge_{(h)}}/p^n$ , it is idempotent and flat, so gives a basic setup in the sense of [GR, 2.1.1]. We thus regard the pair  $(\mathbb{Z}[q^{1/p^\infty}]^{\wedge_{(p,q-1)}}, W^{(p)}(\mathfrak{m}))$  as an inverse system of basic setups for almost ring theory.

We then follow the terminology and notation of [GR], studying  $p$ -adically complete  $(\mathbb{Z}[q^{1/p^\infty}]^{\wedge_{(p,q-1)}})^a$ -modules (almost  $\mathbb{Z}[q^{1/p^\infty}]^{\wedge_{(p,q-1)}}$ -modules) given by localising at almost isomorphisms, the maps whose kernel and cokernel are  $W^{(p)}(\mathfrak{m})$ -torsion.

The obvious functor  $(-)^a$  from modules to almost modules has a right adjoint  $(-)_*$ , given by  $N_* := \mathrm{Hom}_{\mathbb{Z}[q^{1/p^\infty}]^{\wedge_{(p,q-1)}}}(W^{(p)}(\mathfrak{m}), N)$ , the module of almost elements. Since the counit  $(M_*)^a \rightarrow M$  of the adjunction is an (almost) isomorphism, we may also regard almost modules as a full subcategory of the category of modules, consisting of those  $M$  for which the natural map  $M \rightarrow (M^a)_*$  is an isomorphism. We can define  $p$ -adically complete  $(\mathbb{Z}[q^{1/p^\infty}]^{\wedge_{(p,q-1)}})^a$ -algebras similarly, forming a full subcategory of  $\mathbb{Z}[q^{1/p^\infty}]^{\wedge_{(p,q-1)}}$ -algebras.

**Lemma 3.5.** *For any  $\mathbb{Z}[[q-1]]$ -module  $M$ , we may recover the  $\mathbb{Z}[q^{1/p^\infty}]^{\wedge_{(p,q-1)}}$ -module  $(M \otimes_{\mathbb{Z}[q]} \mathbb{Z}[q^{1/p^\infty}])^{\wedge_p}$  as the module of almost elements of the associated almost  $\mathbb{Z}[q^{1/p^\infty}]^{\wedge_{(p,q-1)}}$ -module.*

*Proof.* Since  $M \otimes_{\mathbb{Z}[q]} \mathbb{Z}[q^{1/p^\infty}] = \bigoplus_{\alpha} M \otimes q^\alpha$  for  $\alpha \in p^{-\infty}\mathbb{Z}$  with  $0 \leq \alpha < 1$ , calculation shows that  $(M \otimes_{\mathbb{Z}[q]} \mathbb{Z}[q^{1/p^\infty}])^{\wedge_p} \rightarrow (M \otimes_{\mathbb{Z}[q]} \mathbb{Z}[q^{1/p^\infty}])_*^{\wedge_p}$  must be an isomorphism.  $\square$

**3.3. Perfectoid algebras.** We now relate Scholze's perfectoid algebras to a class of  $\Lambda_p$ -rings.

**Definition 3.6.** Define a perfectoid  $\Lambda_p$ -ring to be a flat  $p$ -adically complete  $\Lambda_p$ -algebra over  $\mathbb{Z}_p$ , on which the Adams operation  $\Psi^p$  is an isomorphism.

For a perfectoid field  $K$  in the sense of [Sch1], there is a tilt  $K^\flat$  (a complete perfect field of characteristic  $p$ ). The subring of power-bounded elements is denoted  $K^\circ \subset K$ .

**Lemma 3.7.** *Given a perfectoid field  $K$ , we have equivalences*

$$\begin{array}{ccc} \text{perfectoid almost } K^\circ\text{-algebras} & & \\ K^{\circ a} \otimes_{(\mathcal{A}_{\text{inf}}(K^\circ)^a)} - & \begin{array}{c} \uparrow \\ \mathcal{A}_{\text{inf}} \\ \downarrow \end{array} & \\ \text{perfectoid almost } \Lambda_p\text{-rings over } \mathcal{A}_{\text{inf}}(K^\circ) & & \\ \mathbb{F}_p \otimes_{\mathbb{Z}_p} - & \begin{array}{c} \downarrow \\ W^{(p)} \\ \uparrow \end{array} & \\ \text{perfectoid almost } K^{\flat o}\text{-algebras} & & \end{array}$$

of categories, where  $\mathcal{A}_{\text{inf}}(C) := \varprojlim_{\Psi^p} W^{(p)}(C)$ .

*Proof.* A perfectoid  $\Lambda_p$ -ring  $B$  is a deformation of the  $\mathbb{F}_p$ -algebra  $B/p$ . As in [Sch1, Proposition 5.13], a perfect  $\mathbb{F}_p$ -algebra  $C$  has a unique deformation  $W^{(p)}C$  over  $\mathbb{Z}_p$ , to which Frobenius must lift uniquely; this gives the bottom pair of equivalences.

We then observe that since  $B := \mathcal{A}_{\text{inf}}(C)$  is a perfectoid  $\Lambda_p$ -ring for any flat  $p$ -adically complete  $\mathbb{Z}_p$ -algebra  $C$ , we must have  $B \cong W^{(p)}(B/p)$ . Comparing rank 1 elements then gives a monoid isomorphism  $(B/p) \cong \varprojlim_{x \mapsto x^p} C$ , from which it follows that

$$\mathbb{F}_p \otimes_{\mathbb{Z}_p} \mathcal{A}_{\text{inf}}(C) \cong \varprojlim_{\Phi} (C/p) = C^\flat$$

whenever  $C$  is perfectoid. Since tilting gives an equivalence of almost algebras by [Sch1, Theorem 5.2], this completes the proof.  $\square$

We will only apply Lemma 3.7 to perfectoid almost  $\Lambda_p$ -rings over  $\mathbb{Z}[q^{1/p^\infty}]^{\wedge_{(p,q-1)}}$ , in which case it shows that reduction modulo  $[p]_{q^{1/p}}$  (resp.  $p$ ) gives an equivalence with perfectoid  $(\mathbb{Z}[\zeta_{p^\infty}]^{\wedge_p})^a$ -algebras (resp. perfectoid  $(\mathbb{F}_p[q^{1/p^\infty}]^{\wedge_{(q-1)}})^a$ -algebras),

**3.4. Functoriality of  $q$ -de Rham cohomology.** Since  $(\Psi^{1/p^\infty} \widehat{\text{qDR}}_p(A/\mathbb{Z}_p))^{\wedge_p}$  is represented by a cosimplicial perfectoid  $\Lambda_p$  ring over  $\mathbb{Z}[q^{1/p^\infty}]^{\wedge_{(p,q-1)}}$  for any flat  $\Lambda_p$ -ring  $A$  over  $\mathbb{Z}_p$ , it corresponds under Lemma 3.7 to a cosimplicial perfectoid  $(\mathbb{Z}[\zeta_{p^\infty}]^{\wedge_p})^a$ -algebra, representing the following functor:

**Lemma 3.8.** *For a perfectoid  $(\mathbb{Z}[\zeta_{p^\infty}]^{\wedge_p})^a$ -algebra  $C$ , and a  $\Lambda_p$ -ring  $A$  over  $\mathbb{Z}_p$  with  $X = \text{Spec } A$ , there is a canonical isomorphism*

$$X_{\text{strat}}^q(\varprojlim_{\Psi^p} W^{(p)}(C)_*) \cong \text{Im}(\varprojlim_{\Psi^p} X(C_*) \rightarrow X(C_*)),$$

for the ring  $C_*$  of almost elements.

*Proof.* By definition,  $X_{\text{strat}}^q(\varprojlim_{\Psi^p} W^{(p)}(C)_*)$  is the image of

$$\text{Hom}_{\Lambda_p}(A, \varprojlim_{\Psi^p} W^{(p)}(C)_*) \rightarrow \text{Hom}_{\Lambda_p}(A, (\varprojlim_{\Psi^p} W^{(p)}(C)_*)/(q-1)).$$

Since right adjoints commute with limits, we may rewrite the first term as  $\varprojlim_{\Psi^p} \mathrm{Hom}_{\Lambda_p}(A, W^{(p)}(C_*)) = \varprojlim_{\Psi^p} X(C_*)$ .

Setting  $B := \varprojlim_{\Psi^p} W^{(p)}(C)_*$ , observe that because  $[p^n]_{q^{1/p^n}}(q^{1/p^n} - 1) = (q - 1)$ , we have  $\bigcap_n [p^n]_{q^{1/p^n}} B = (q - 1)B$ , any element on the left defining an almost element of  $(q - 1)B$ , hence a genuine element since  $B = B_*$  is flat. Then note that since the projection map  $\theta: B \rightarrow C_*$  has kernel  $([p]_{q^{1/p}})$ , the map  $\theta \circ \Psi^{p^{n-1}}$  has kernel  $([p]_{q^{1/p^n}})$ , and so  $B \rightarrow W^{(p)}C_*$  has kernel  $\bigcap_n [p^n]_{q^{1/p^n}} B$ . Thus

$$\mathrm{Hom}_{\Lambda_p}(A, (\varprojlim_{\Psi^p} W^{(p)}(C)_*)/(q - 1) \hookrightarrow \mathrm{Hom}_{\Lambda_p}(A, W^{(p)}C_*) = X(C_*).$$

□

In fact, the tilting equivalence gives  $\varprojlim_{\Psi^p} X(C_*) \cong X(C_*^\flat)$ , so the only dependence of  $((\Psi^{1/p^\infty} \widehat{\mathrm{qDR}}_p(A/\mathbb{Z}_p))^{\wedge p})^a$  on the Frobenius lift  $\Psi^p$  is in determining the image of  $X(C_*^\flat) \rightarrow X(C_*)$ .

Although this map is not surjective, it is almost so in a precise sense, which we now use to establish independence of  $\Psi^p$ , showing that, up to faithfully flat descent,  $\mathrm{q}\hat{\mathrm{DR}}_p(A/\mathbb{Z}_p)^{\wedge p}/[p]_{q^{1/p}}$  is the best possible perfectoid approximation to  $A[\zeta_{p^\infty}]^{\wedge p}$ .

**Definition 3.9.** Given a functor  $X$  from  $(\mathbb{Z}[\zeta_{p^\infty}]^{\wedge p})^a$ -algebras to sets and a functor  $\mathcal{A}$  from perfectoid  $(\mathbb{Z}[\zeta_{p^\infty}]^{\wedge p})^a$ -algebras to abelian groups, we write

$$\mathbf{R}\Gamma_{\mathrm{Pfd}}(X, \mathcal{A}) := \mathbf{R}\mathrm{Hom}_{[\mathrm{Pfd}((\mathbb{Z}_p[\zeta_{p^\infty}]^{\wedge p})^a), \mathrm{Set}]}(X, \mathcal{A}),$$

where  $\mathrm{Pfd}(S^a)$  denotes the category of perfectoid almost  $S$ -algebras, and  $[\mathcal{C}, \mathrm{Set}]$  denotes set-valued functors on  $\mathcal{C}$ . When  $X$  is representable by a  $(\mathbb{Z}[\zeta_{p^\infty}]^{\wedge p})^a$ -algebra  $C$ , we simply denote this by  $\mathbf{R}\Gamma_{\mathrm{Pfd}}(C, \mathcal{A})$  — when  $C$  is perfectoid, this will just be  $\mathcal{A}(C)$ .

The following gives a refinement of [BMS, Theorem 1.17], addressing some of the questions in [BMS, Remark 1.11]:

**Theorem 3.10.** *If  $R$  is a  $p$ -adically complete  $\Lambda_p$ -ring over  $\mathbb{Z}_p$ , and  $A$  a formal  $R$ -deformation of a smooth ring over  $(R/p)$ , then the complex*

$$\mathbf{R}\Gamma_{\mathrm{Pfd}}((A[\zeta_{p^\infty}] \otimes_R \Psi^{1/p^\infty} R)^{\wedge p}, \mathcal{A}_{\mathrm{inf}})$$

*of  $\mathbb{Z}_p[q^{1/p^\infty}]^{\wedge(p, q-1)}$ -modules is almost quasi-isomorphic to  $(\Psi^{1/p^\infty} \mathrm{q}\hat{\mathrm{DR}}_p(A/R))^{\wedge p}$  for any  $\Lambda_p$ -ring structure on  $A$  coming from a framing over  $R$  as in Theorem 2.7.*

*Proof.* First observe that  $((\Psi^{1/p^\infty} \mathrm{q}\hat{\mathrm{DR}}_p(A/R))^{\wedge p})^a$  is the completion of  $\mathrm{q}\hat{\mathrm{DR}}_p(A/R)$  with respect to the category of cosimplicial perfectoid almost  $\Lambda_p$ -rings over  $\mathbb{Z}[q^{1/p^\infty}]^{\wedge(p, q-1)}$ . Combining the definition of  $\widehat{\mathrm{qDR}}_p$  with Lemma 3.7, it then follows that for  $X = \mathrm{Spec} A$  and  $Y = \mathrm{Spec} R$ , the complex  $(\widehat{\mathrm{qDR}}_p(A/R)^{\wedge p})_*$  is given by the homotopy limit

$$\mathbf{R}\Gamma_{\mathrm{Pfd}}((X_{\mathrm{strat}}^q \times_{Y_{\mathrm{strat}}^q} Y) \circ (\mathcal{A}_{\mathrm{inf}})_*, (\mathcal{A}_{\mathrm{inf}})_*).$$

Writing  $X^\infty(C) := \mathrm{Im}(\varprojlim_{\Psi^p} X(C_*) \rightarrow X(C_*))$ , Lemma 3.8 then combines with the description above to give

$$\begin{aligned} (\widehat{\mathrm{qDR}}_p(A/R)^{\wedge p})_* &\simeq \mathbf{R}\Gamma_{\mathrm{Pfd}}(X^\infty \times_{Y^\infty} \varprojlim_{\Psi^p} Y, (\mathcal{A}_{\mathrm{inf}})_*), \\ &\simeq \mathbf{R}\Gamma_{\mathrm{Pfd}}(X^\infty \times_Y \varprojlim_{\Psi^p} Y, (\mathcal{A}_{\mathrm{inf}})_*). \end{aligned}$$



We now introduce a Grothendieck topology on the category  $[\mathrm{Pfd}_{(\mathbb{Z}[\zeta_{p^\infty}]^{\wedge p})^a}, \mathrm{Set}]$  by taking covering morphisms to be those maps  $C \rightarrow C'$  of perfectoid algebras which are almost faithfully flat modulo  $p$ . Since  $C^\flat = \varprojlim_{\Phi} (C/p)$ , the functor  $\mathcal{A}_{\mathrm{inf}}$  satisfies descent with respect to these coverings, so the map

$$\mathbf{R}\Gamma_{\mathrm{Pfd}}((X^\infty \times_Y \varprojlim_{\Psi^p} Y)^\sharp, (\mathcal{A}_{\mathrm{inf}})_*) \rightarrow \mathbf{R}\Gamma_{\mathrm{Pfd}}(X^\infty \times_Y \varprojlim_{\Psi^p} Y, (\mathcal{A}_{\mathrm{inf}})_*)$$

is a quasi-isomorphism, where  $(-)^{\sharp}$  denotes sheafification.

In other words, the calculation of  $(\widehat{\mathrm{qDR}}_p(A/R)^{\wedge p})^a$  is not affected if we tweak the definition of  $X^\infty$  by taking the image sheaf instead of the image presheaf. We then have

$$(X^\infty)^\sharp(C) = \bigcup_{C \rightarrow C'} \mathrm{Im}(X(C_*) \times_{X(C'_*)} \varprojlim_{\Psi^p} X(C'_*) \rightarrow X(C_*)),$$

where  $C \rightarrow C'$  runs over all covering morphisms.

Now,  $\varprojlim_{\Psi^p} X$  is represented by the perfectoid algebra  $(\Psi^{1/p^\infty} A)^{\wedge p}$ , which is isomorphic to  $A[x_1^{1/p^\infty}, \dots, x_d^{1/p^\infty}]^{\wedge p}$  as in the proof of Lemma 3.3. This allows us to appeal to André's results [And, §2.5] as generalised in [Bha, Theorem 2.3]. For any morphism  $f: A \rightarrow C$ , there exists a covering morphism  $C \rightarrow C_i$  such that  $f(x_i)$  has arbitrary  $p$ -power roots in  $C_i$ . Setting  $C' := C_1 \otimes_C \dots \otimes_C C_d$ , this means that the composite  $A \xrightarrow{f} C \rightarrow C'$  extends to a map  $(\Psi^{1/p^\infty} A)^{\wedge p} \rightarrow C'$ , so  $f \in (X^\infty)^\sharp(C)$ . We have thus shown that  $(X^\infty)^\sharp = X$ , giving the required equivalence

$$((\Psi^{1/p^\infty} \widehat{\mathrm{qDR}}_p(A/R))^{\wedge p})_* \simeq \mathbf{R}\Gamma_{\mathrm{Pfd}}(X \times_Y \varprojlim_{\Psi^p} Y, (\mathcal{A}_{\mathrm{inf}})_*).$$

□

**Corollary 3.11.** *If  $R$  is a  $p$ -adically complete  $\Lambda_p$ -ring over  $\mathbb{Z}_p$ , and  $A$  a formal  $R$ -deformation of a smooth ring over  $(R/p)$ , then the  $q$ -de Rham cohomology complex  $(q\text{-}\Omega_{A/R, \square}^\bullet \otimes_{R[q]} (\Psi^{1/p^\infty} R)[q^{1/p^\infty}])^{\wedge p}$  is, up to quasi-isomorphism, independent of a choice of co-ordinates □. It is naturally an invariant of the commutative  $p$ -adically complete  $(\Psi^{1/p^\infty} R)[\zeta_{p^\infty}]^{\wedge p}$ -algebra  $(A[\zeta_{p^\infty}] \otimes_R \Psi^{1/p^\infty} R)^{\wedge p}$ .*

*Proof.* By Theorem 3.10, we know that the complex  $((\Psi^{1/p^\infty} \widehat{\mathrm{qDR}}_p(A/R))^{\wedge p})_*$  depends only on  $(A[\zeta_{p^\infty}] \otimes_R \Psi^{1/p^\infty} R)^{\wedge p}$ . Since

$$\Psi^{1/p^\infty} \widehat{\mathrm{qDR}}_p(A/R) = \Psi^{1/p^\infty} \widehat{\mathrm{qDR}}_p((A \otimes_R \Psi^{1/p^\infty} R)/\Psi^{1/p^\infty} R),$$

Theorem 2.7 combines with Lemmas 3.3 and 3.5 to give

$$(q\text{-}\Omega_{A/R, \square}^\bullet \otimes_{R[q]} (\Psi^{1/p^\infty} R)[q^{1/p^\infty}])^{\wedge p} \simeq \mathbf{L}\eta_{(q-1)}((\Psi^{1/p^\infty} \widehat{\mathrm{qDR}}_p(A/R))^{\wedge p})_*,$$

which completes the proof. □

*Remark 3.12* (Scholze's conjectures). If we weaken the conjectures of [Sch2] by attaching roots of  $q$ , then most of them follow from the results above. Taking  $R = \mathbb{Z}_p$  in Corollary 3.11 establishes the analogue of [Sch2, Conjecture 3.1] (co-ordinate independence of  $q$ -de Rham cohomology over  $\mathbb{Z}$ ) via an arithmetic fracture square. Taking more general base rings  $R$  in Corollary 3.11 gives an analogue of [Sch2, Conjecture 7.1], further weakened by having to invert all Adams operations on  $R$ .

The description of Theorem 3.10 is very closely related to the definition of  $A\Omega^\bullet$  in [BMS, Definition 1.12], giving an analogue of [Sch2, Conjecture 4.3], and hence the comparison with singular cohomology in [Sch2, Conjecture 3.3]. The operations

described in [Sch2, Conjectures 6.1 and 6.2] correspond to the Adams operations on  $q$ DR respectively at and away from the residue characteristics. Remark 1.19 provides a category of  $q$ -connections as described in [Sch2, Conjecture 7.5]; these will correspond via Theorem 1.11 to projective  $\mathcal{A}_{\text{inf}}$ -modules on the site of integral perfectoid algebras over  $A[\zeta_{p^\infty}]^{\wedge p}$ , so are again independent of co-ordinates after base change to  $\mathbb{Z}[q^{1/p^\infty}]$ .

## REFERENCES

- [And] Y. Andre. La conjecture du facteur direct. arXiv: 1609.00345 [math.AG], 2016.
- [Bha] B. Bhatt. On the direct summand conjecture and its derived variant. arXiv:1608.08882[math.AG], 2016.
- [BMS] B. Bhatt, M. Morrow, and P. Scholze. Integral  $p$ -adic Hodge theory. arXiv:1602.03148 [math.NT], 2016.
- [Bor] James Borger. The basic geometry of Witt vectors, I: The affine case. *Algebra Number Theory*, 5(2):231–285, 2011.
- [GR] Ofer Gabber and Lorenzo Ramero. *Almost ring theory*, volume 1800 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2003. arXiv:math/0201175v3 [math.AG].
- [Gro] A. Grothendieck. Crystals and the de Rham cohomology of schemes. In *Dix Exposés sur la Cohomologie des Schémas*, pages 306–358. North-Holland, Amsterdam, 1968.
- [Joy] André Joyal.  $\delta$ -anneaux et vecteurs de Witt. *C. R. Math. Rep. Acad. Sci. Canada*, 7(3):177–182, 1985.
- [Sch1] Peter Scholze. Perfectoid spaces. *Publ. Math. Inst. Hautes Études Sci.*, 116:245–313, 2012.
- [Sch2] Peter Scholze. Canonical  $q$ -deformations in arithmetic geometry. *Ann. Fac. Sci. Toulouse Math.*, to appear. arXiv:1606.01796v1[math.AG].
- [Sim] Carlos Simpson. Homotopy over the complex numbers and generalized de Rham cohomology. In *Moduli of vector bundles (Sanda, 1994; Kyoto, 1994)*, volume 179 of *Lecture Notes in Pure and Appl. Math.*, pages 229–263. Dekker, New York, 1996.